rst Aic **INFANTS AND CHILDREN**

To handle an emergency situation, use Emergency Scene Management (ESM).

- 1 Take charge.
- 2 Call out for help.
- Assess hazards and make the area safe.
- Find out what happened.

- 5 Identify yourself and offer to help.
- If head or spinal injuries are suspected, support the head and neck.
- Assess responsiveness.
- 8 Send or go for medical help.

Note: Protect yourself and others by wearing non-latex gloves when giving first aid. Use a shield or face mask with a one-way valve when giving CPR.

Cardiopulmonary Resuscitation (CPR)

- Open airway push back on forehead and lift chin.
- Check breathing. If the casualty is not breathing...
- Make a tight seal over mouth and nose of an infant (under 1 year). Pinch nose of a child (1-8 years) and make a tight seal over the mouth. Give 2 breaths.
- Make sure casualty is on a firm flat surface.
- For a child place hands on centre of chest
- Position shoulders directly over hands and keep elbows locked.
- Compress firmly 30 times then give 2 breaths. Continue cycles of 30 compressions and 2 breaths until help arrives.



CHECK FOR BREATHING



BREATHE INTO CASUALTY 2X



PLACE HANDS ON CENTRE





COMPRESS FIRMLY 30 TIMES THEN GIVE 2 BREATHS. CONTINUE CYCLES OF 30 COMPRESSIONS AND 2 BREATHS UNTIL HELP ARRIVES.

- For an Infant place 2 fingertips on the infant's breastbone just below the nipples.
- Use 2 fingers to press down firmly on the breastbone.
- Compress firmly 30 times then give 2 breaths.
- Continue cycles of 30 compressions and 2 breaths until help arrives.







PRESS DOWN FIRMLY 30 TIMES THEN GIVE 2 BREATHS... REPEAT UNTIL HELP ARRIVES


Choking (Child)

If a choking child can speak, breathe or cough - STAND BY and encourage coughing.

If a conscious child cannot speak, breathe or cough:

- Stand behind the child and find top of the hip bones with your hands.
- Place a fist midline against the abdomen.
- Grasp fist with other hand and press inward and upward forcefully.
- Continue until object is expelled or child becomes unconscious.

If the child becomes unconscious ease him or her to the ground and send for medical help



- Open the airway and check breathing. If not breathing...
- Give two breaths. If air doesn't go in the first time, reposition the head and try again.
- Begin CPR. Give 30 compressions.
- Each time you finish 30 compressions, look in mouth before giving a breath.



PLACE FIST MID-



Choking (Infant)

If infant is making high pitched noises or is having trouble breathing:

- Give 5 forceful blows between the shoulder blades.
- Turn infant over and give five chest thrusts using two fingers on the infant's breastbone just below the nipple line.
- Repeat 5 back blows and 5 chest thrusts.
- Continue until object is expelled or child becomes unconscious.

If the infant becomes unconscious place him or her on a firm flat surface and send for medical help

- Open the mouth and look for obstruction.
- Open the airway and check breathing. If not breathing...
- Give two breaths. If air doesn't go in the first time, reposition the head and try again.
- Begin CPR. Give 30 compressions.
 - Each time you finish 30 compressions, look in mouth before giving a breath.



Bleeding To control severe bleeding

Immediately apply direct pressure to the wound over a pad of dressings. Keep the casualty lying down.



CONTROL BLEEDING IMMEDIATELY

Unconsciousness

Get medical help. Make certain child is breathing and then place in the recovery position.

If not breathing begin CPR



RECOVERY POSITIO

AMBULANCE FIRE POLICE FAMILY DOCTOR **POISON INFORMATION CENTRE**

NOTE: This poster is a step-by-step guide to what you can do until medical help arrives. These tips do not take the place of first aid training.

